

行政院國家科學委員會專題研究計劃成果報告

* 行進波之函數理論法 *

* *

計劃類別：☒ 個別計劃 ☐ 整合型計劃

計劃編號：NSC 90-2115-M-032-005

執行期間：90 年 8 月 1 日至 91 年 7 月 31 日

個別型計劃：計劃主持人：張慧京
國科會助理：喻芝健
國科會助理：劉建德
共同主持人：

整合型計劃：總計劃主持人：
子計劃主持人：

註：整合型計劃總報告與子計劃成果報告請分開編印各成一冊
，彙整一起繳送國科會。

處理方式：☐ 可立即對外提供參考(請打√) ☒ 一年後可對外提供參考☐ 兩年後可對外提供參考

(必要時，本會得展延發表時限)

執行單位：淡江大學數學系

中華民國 91 年 10 月 25 日

中文摘要:

在此報告中我們研究由無限個蔡氏電路耦合所形成的理想化的偏微分方程之行進波解的存在性。我們先利用變數變換的方法將原偏微分方程轉換成一邊界值問題，再利用上近似解與下近似解的方法得到當參數在某些值域時此邊界值微分方程有異源解。因而我們可以得到耦合蔡式電路有任意速度的行進波解的結論。

關鍵詞：

上近似解，下近似解，Nagumo 條件，行進波解，異源解。

英文摘要:

In this work, we study the existence of heteroclinic orbits for a ordinary differential equations which arise from a one-dimensional array of Chua's circuits. By using the upper and lower solutions method and a zero-order approximation, we show that for a certain set of parameters we have the existence of traveling wave solutions for the idealized system for some given speed.

關鍵詞：

upper solution, lower solution, Nagumo condition, traveling solution, heteroclinic orbit.

1 Summary

In this work we will study the partial differential equation which had been derived by idealizing the infinitely coupled chua's circuits.

$$\begin{aligned}\frac{\partial U}{\partial t} &= \alpha(Z - f(U)) + D\frac{\partial^2 U}{\partial x^2}, \quad D > 0, \\ \frac{\partial Z}{\partial t} &= U - Z + W, \\ \frac{\partial W}{\partial t} &= -\beta Z.\end{aligned}\tag{1.1}$$

where

$$f(u) = \begin{cases} m_0 u, & u \leq \bar{u}_1, \\ -m_1(u - \bar{u}), & \bar{u}_1 < u \leq \bar{u}_2, \\ m_2(u - 1), & u > \bar{u}_2. \end{cases}$$

Mainly, we are interesting in the existence of traveling solution of equation (1.1). Let c be the wave speed of the traveling wave solution. By introducing a moving coordinate $t' = \frac{t'}{-c} = \frac{1}{-c}(x - ct)$ and setting

$$U(t, x) = u\left(\frac{x}{-c} + t\right), \quad Z(t, x) = z\left(\frac{x}{-c} + t\right), \quad W(t, x) = w\left(\frac{x}{-c} + t\right),$$

we arrive at ordinary differential equations for (u, z, w) :

$$\begin{cases} \varepsilon \ddot{u} = \dot{u} - \alpha(z - f(u)), \\ \dot{z} = u - z + w, \\ \dot{w} = -\beta z \end{cases}\tag{1.2}$$

with the boundary conditions

$$\lim_{t \rightarrow -\infty} (u, \dot{u}, z, w) = (0, 0, 0, 0), \quad \lim_{t \rightarrow +\infty} (u, \dot{u}, z, w) = (1, 0, 0, -1), \tag{1.3}$$

where $\varepsilon = \frac{D}{c^2}$.

To show that the system (1.2) with boundary condition (1.3) has a solution for a certain set of parameter values. We will need the following existence theorems for a certain class of boundary value problem.

Let $\mathcal{T} : C^0([a, b], \mathbb{R}) \rightarrow C^2([a, b], \mathbb{R})$ be a bounded linear operator and $f : [a, b] \times \mathbb{R} \times \mathbb{R} \times C^0([a, b], \mathbb{R}) \rightarrow \mathbb{R}$ be continuous.

Theorem 1.1: If $f(t, y, z, [\mathcal{T}\xi])$ is bounded on $[a, b] \times \mathbb{R} \times \mathbb{R} \times C^1([a, b], \mathbb{R})$. Then for and $A, B \in \mathbb{R}$, the boundary value problem

$$\begin{aligned} y'' &= f(t, y, y', [\mathcal{T}y]), \\ y(a) &= A, \quad y(b) = B \end{aligned}$$

has a solution.

Definition 1.2: Let $\underline{\omega}$ and $\bar{\omega}$ be continuous and piecewise C^2 functions with $\underline{\omega}(t) \leq \bar{\omega}(t)$ on $[a, b]$. That is, there is a finite partition $\{t_i\}$, $i = 0, \dots, n$, of $[a, b]$ with $a = t_0 < t_1 < t_2 < \dots < t_n = b$, such that on each closed subinterval $[t_{i-1}, t_i]$, $\underline{\omega}$ and $\bar{\omega}$ are twice continuously differentiable. At the partition points t_{i-1} and t_i , the right and left-handed derivatives satisfy the following:

$$\underline{\omega}'(t_i^-) < \underline{\omega}'(t_i^+), \quad \bar{\omega}'(t_i^-) > \bar{\omega}'(t_i^+), \quad i = 1, 2, \dots, n-1,$$

In addition, if on each open subinterval (t_{i-1}, t_i) the following inequalities are satisfied for any $\xi(t) \in C^0([a, b])$ satisfying $\underline{\omega}(t) \leq \xi(t) \leq \bar{\omega}(t)$,

$$\begin{aligned} \underline{\omega}''(t) &\geq f(t, \underline{\omega}(t), \underline{\omega}'(t), [\mathcal{T}\xi](t)), \\ \bar{\omega}''(t) &\leq f(t, \bar{\omega}(t), \bar{\omega}'(t), [\mathcal{T}\xi](t)) \end{aligned}$$

then $\underline{\omega}(t)$ and $\bar{\omega}(t)$ are called the piecewise C^2 lower and upper solution, respectively, of the differential equation

$$y'' = f(t, y, y', [\mathcal{T}\xi(t)]) \tag{1.4}$$

Definition 1.3: Let $\underline{\omega}, \bar{\omega} \in C^0([a, b], \mathbb{R})$ with $\underline{\omega}(t) \leq \bar{\omega}(t)$ on $[a, b]$ and $h \in C^0(\mathbb{R}^+, \mathbb{R}^+)$ satisfy

$$\int_{\lambda}^{\infty} \frac{s ds}{h(s)} > \max_{t \in [a, b]} \bar{\omega}(t) - \min_{t \in [a, b]} \underline{\omega}(t),$$

where

$$\lambda(b - a) = \max\{|\underline{\omega}(a) - \bar{\omega}(b)|, |\underline{\omega}(b) - \bar{\omega}(a)|\}.$$

The differential equation (1.4) is said to satisfy a Nagumo condition on $[a, b]$ with respect to the pair $\underline{\omega}, \bar{\omega}$ if the following inequality is satisfied for any $t \in [a, b]$, $\underline{\omega}(t) \leq u \leq \bar{\omega}(t)$, $v \in \mathbb{R}$ and any $\xi \in C^0([a, b], \mathbb{R})$ with $\underline{\omega}(t) \leq \xi(t) \leq \bar{\omega}(t)$,

$$|f(t, u, v, [\mathcal{T}\xi](t))| \leq h(|v|).$$

Theorem 1.4: Suppose that $\underline{\omega}$ and $\bar{\omega}$ are piecewise C^2 lower and upper solutions of the differential equation

$$y'' = f(t, y, y', [\mathcal{T}\xi](t)),$$

where $f : [a, b] \times \mathbb{R} \times \mathbb{R} \times C^0([a, b], \mathbb{R})$ is continuous and $\mathcal{T} : C^0([a, b], \mathbb{R}) \rightarrow C^2([a, b], \mathbb{R})$ is a bounded linear operator. Assume that $y'' = f(t, y, y', [\mathcal{T}\xi](t))$ satisfies a Nagumo condition on $[a, b]$ with respect to the pair $\underline{\omega}$ and $\bar{\omega}$. Then for any A, B with $\underline{\omega}(a) \leq A \leq \bar{\omega}(a)$ and $\underline{\omega}(b) \leq B \leq \bar{\omega}(b)$, the following boundary value problem

$$y'' = f(t, y, y', [\mathcal{T}y]),$$

$$y(a) = A, \quad y(b) = B,$$

has a solution $y(t)$ with $\underline{\omega}(t) \leq y(t) \leq \bar{\omega}(t)$ on $[a, b]$. Moreover, let $N_0 > 0$ be determined by:

$$\int_{\lambda}^{N_0} \frac{s ds}{h(s)} = \max_{t \in [a, b]} \bar{\omega}(t) - \min_{t \in [a, b]} \underline{\omega}(t),$$

then $|y'(t)| \leq N_0$, for all $a \leq t \leq b$.

Now let's define the zero order approximation solution (u_0, z_0, w_0) of (1.2) as follows

$$\begin{aligned} u_0(t) &= \begin{cases} u_0^L(t) = e^{\frac{t}{\varepsilon}}, & t \leq 0, \\ u_0^R(t) = 1, & t > 0. \end{cases} \\ z_0(t) &= \begin{cases} z_0^L(t) = \frac{\varepsilon}{1+\varepsilon+\beta\varepsilon^2} e^{\frac{t}{\varepsilon}}, & t \leq 0, \\ z_0^R(t) = [\frac{\varepsilon}{1+\varepsilon+\beta\varepsilon^2} \cos(\bar{\beta}t) + (\frac{1}{\bar{\beta}} - \frac{(1+2\beta\varepsilon)\varepsilon}{\bar{\beta}(1+\varepsilon+\beta\varepsilon^2)}) \sin(\bar{\beta}t)] e^{-\frac{1}{2}t}, & t > 0, \end{cases} \\ w_0(t) &= \begin{cases} w_0^L(t) = -\frac{\beta\varepsilon^2}{1+\varepsilon+\beta\varepsilon^2} e^{\frac{t}{\varepsilon}}, & t \leq 0, \\ w_0^R(t) = [\frac{1+\varepsilon}{1+\varepsilon+\beta\varepsilon^2} \cos(\bar{\beta}t) + \frac{1+(1-2\beta)\varepsilon}{2\bar{\beta}(1+\varepsilon+\beta\varepsilon^2)} \sin(\bar{\beta}t)] e^{-\frac{1}{2}t} - 1, & t > 0, \end{cases} \end{aligned}$$

where $\bar{\beta} = \sqrt{\beta - \frac{1}{4}}$.

Our main result is as follows.

Theorem 1.5. Consider the following differential equation

$$\begin{cases} \varepsilon \ddot{u} = \dot{u} - \alpha(z - f(u)), \\ \dot{z} = u - z + w, \\ \dot{w} = -\beta z \end{cases}$$

with the boundary conditions

$$\lim_{t \rightarrow -\infty} (u, \dot{u}, z, w) = (0, 0, 0, 0), \quad \lim_{t \rightarrow +\infty} (u, \dot{u}, z, w) = (1, 0, 0, -1),$$

If $\varepsilon > 0$ is sufficiently small, $\beta > \frac{1}{4}$ and $a > 0$ is sufficiently large, then for any $0 < b < \frac{1}{2}$ the boundary value problem (1.2) (1.3) has solution (u, \dot{u}, z, w) .

Moreover, they satisfy the following estimates

$$\begin{aligned} |(u, z, w) - (u_0^L(t), z_0^L(t), w_0^L(t))| &\leq Me^{at}, & t \leq 0, \\ |(u, z, w) - (u_0^R(t), z_0^R(t), w_0^R(t))| &\leq Me^{-bt}, & t > 0. \end{aligned}$$

where some appropriate constant M .

References

- Bernfeld, S. R. & Lakshmikantham, V. [1974] *An introduction to Nonlinear Boundary Value Problems* (Academic Press, New York)
- Chow, S.-N., Mallet-Paret, J. & Shen, W. [1998] “ Traveling wave in lattice dynamical systems,” *J. Differential Equations* **149** , 248-291.
- Chua, L. O. [1992] “ The genesis of Chua’s circuit,” *Archiv für Elektronik und Übertragungstechnik* **46**, 250-257.
- Chua, L. O. [1998] “ CNN: A paradigm for complexity,” *World Scientific Series on Nonlinear Science*, ” Series A - Vol. 31, (World Scientific, Singapore)
- De Coster, C. & Habets, P. [1996] “ Upper and Lower Solution in the Theory of ODE Boundary Value Problem: Classical and Recent Results,” in *Nonlinear Analysis and BVP for ODE* , eds. Zanolin, F.(Springer Wien , New York) **No. 371**, pp. 1-78.
- Perez-Munuzuri, V., Perez-Villar, V. & Chua, L. O. [1992] “ Propagation failure in linear arrays of Chua’s circuits,” *Int. J. Bifur. Chaos* **2**, 403-406.
- Perez-Munuzuri, V., Perez-Villar, V. & Chua, L. O. [1993] “ Traveling wave front and its failure in a one-dimensional array of Chua’s circuit,” in *Chua’s circuit: A paradigm for Chaos*, (Word Scientific, Singapore) pp 336-350.